

# 数学物理方程

## 第一部分 2022 年真题解析

1. 针对本征值问题，证明正交性

$$(1). \begin{cases} X''(x) + \lambda X(x) = 0 \\ X'(0) = 0, \quad X'(l) = 0 \end{cases}$$

$$(2). \begin{cases} X''(x) + \lambda X(x) = 0 \\ X(0)\cos\alpha - X'(0)\sin\alpha = 0 \\ X(l)\cos\beta + X'(l)\sin\beta = 0 \end{cases}$$

证明：(1)对于  $X_1(x)$  和  $X_2(x)$  显然有

$$\begin{cases} X_1''(x) + \lambda_1 X_1(x) = 0 \\ X_2''(x) + \lambda_2 X_2(x) = 0 \end{cases}$$

第一个方程乘  $X_2(x)$  减去第二个方程乘  $X_1(x)$  可得

$$X_2(x)X_1''(x) - X_1(x)X_2''(x) = (\lambda_2 - \lambda_1)X_1(x)X_2(x)$$

积分可得

$$\int_0^l [X_2(x)X_1''(x) - X_1(x)X_2''(x)] dx = (\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx$$

$$[X_2(x)X_1'(x) - X_1(x)X_2'(x)] \Big|_0^l = (\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx$$

$$X_2(l)X_1'(l) - X_1(l)X_2'(l) - X_2(0)X_1'(0) + X_1(0)X_2'(0) = (\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx$$

带入边界条件可得

$$(\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx = 0 \xrightarrow{\lambda_2 \neq \lambda_1} \int_0^l X_1(x)X_2(x) dx = 0. \square$$

(2) 同(1)可得

$$X_2(l)X_1'(l) - X_1(l)X_2'(l) - X_2(0)X_1'(0) + X_1(0)X_2'(0) = (\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx$$

$$- X_2'(l)X_1(l)\tan\beta + X_1'(l)X_2(l)\tan\beta - X_2'(0)X_1'(0)\tan\alpha + X_1'(0)X_2'(0)\tan\alpha = (\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx$$

带入边界条件可得

$$(\lambda_2 - \lambda_1) \int_0^l X_1(x)X_2(x) dx = 0 \xrightarrow{\lambda_2 \neq \lambda_1} \int_0^l X_1(x)X_2(x) dx = 0. \square$$

2. 分离变量法求解问题

$$\frac{\partial^2 u}{\partial t^2} - a^2 \frac{\partial^2 u}{\partial x^2} = 9$$

$$u|_{x=0} = 0, \quad u|_{x=l} = 0, \quad u|_{t=0} = 0, \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = 0$$

解：设  $u(x, t) = v(x) + w(x, t)$ ，则有

$$-a^2 v''(x) = 9$$

$$v|_{x=0} = 0, \quad v|_{x=l} = 0$$

求解该方程可得

$$v(x) = -\frac{9}{2a^2}x^2 + \frac{9l}{2a^2}x = \frac{9x(l-x)}{2a^2}$$

而又有

$$\frac{\partial^2 w}{\partial t^2} - a^2 \frac{\partial^2 w}{\partial x^2} = 0$$

$$w|_{x=0} = 0, \quad w|_{x=l} = 0, \quad w|_{t=0} = -v|_{t=0} = \frac{9x(x-l)}{2a^2}, \quad \left. \frac{\partial w}{\partial t} \right|_{t=0} = -\left. \frac{\partial v}{\partial t} \right|_{t=0} = 0$$

带入一般解有

$$w(x, t) = \sum_{n=1}^{\infty} \left( A_n \sin \frac{n\pi a}{l} t + B_n \cos \frac{n\pi a}{l} t \right) \sin \frac{n\pi}{l} x$$

带入边界条件可得

$$A_n = 0 \quad B_n = -\frac{36l^2}{(2n+1)^3 a^2 \pi^3}$$

则有

$$w(x, t) = -\frac{36l^2}{a^2 \pi^3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^3} \cos \frac{(2n+1)\pi a}{l} t \sin \frac{(2n+1)\pi}{l} x$$

综上可得

$$u(x, t) = \frac{9x(l-x)}{2a^2} - \frac{36l^2}{a^2 \pi^3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^3} \cos \frac{(2n+1)\pi a}{l} t \sin \frac{(2n+1)\pi}{l} x. \square$$

注：

$$\begin{aligned} B_n &= \frac{2}{l} \int_0^l \frac{9x(x-l)}{2a^2} \sin \frac{n\pi}{l} x \, dx = \frac{9}{a^2 l} \int_0^l x(x-l) \sin \frac{n\pi}{l} x \, dx \\ &= \frac{9}{a^2 l} \int_0^l x(x-l) \sin \frac{n\pi}{l} x \, dx = \frac{18l^2(\cos n\pi - 1)}{n^3 a^2 \pi^3} = -\frac{36l^2}{(2n+1)^3 a^2 \pi^3} \end{aligned}$$

3. 讨论下列方程类型并化为标准形式

$$y \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

解：判别式： $\Delta = b^2 - ac = -y$ ，则有

(1)  $\Delta < 0 \iff y > 0$ : 该方程属于椭圆型

(2)  $\Delta > 0 \iff y < 0$ : 该方程属于双曲型

(3)  $\Delta = 0 \iff y = 0$ : 该方程属于抛物型

(这里仅仅讨论第一种情况) 特征线:  $\frac{dy}{dx} = \frac{b \pm \sqrt{b^2 - ac}}{a} = \pm \frac{\sqrt{-y}}{y} = \pm i \frac{1}{\sqrt{y}}$ , 解之可得

$$x \pm i \frac{2}{3} y^{\frac{3}{2}} = C$$

作两次换元可得

$$\begin{cases} \xi = x + i \frac{2}{3} y^{\frac{3}{2}} \\ \eta = x - i \frac{2}{3} y^{\frac{3}{2}} \end{cases} \quad \begin{cases} \rho = \xi + \eta = 2x \\ \sigma = i(\xi - \eta) = -\frac{4}{3} y^{\frac{3}{2}} \end{cases}$$

求解可得

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \rho} \frac{\partial \rho}{\partial x} + \frac{\partial u}{\partial \sigma} \frac{\partial \sigma}{\partial x} = 2 \frac{\partial u}{\partial \rho}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left( 2 \frac{\partial u}{\partial \rho} \right) = \frac{\partial}{\partial \rho} \left( 2 \frac{\partial u}{\partial \rho} \right) \frac{\partial \rho}{\partial x} + \frac{\partial}{\partial \sigma} \left( 2 \frac{\partial u}{\partial \rho} \right) \frac{\partial \sigma}{\partial x} = 4 \frac{\partial^2 u}{\partial \rho^2}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial \rho} \frac{\partial \rho}{\partial y} + \frac{\partial u}{\partial \sigma} \frac{\partial \sigma}{\partial y} = -2\sqrt{y} \frac{\partial u}{\partial \sigma}$$

$$\frac{\partial^2 u}{\partial y^2} = \frac{\partial}{\partial y} \left( -2\sqrt{y} \frac{\partial u}{\partial \sigma} \right) = -\frac{1}{\sqrt{y}} \frac{\partial u}{\partial \sigma} - 2\sqrt{y} \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial \sigma} \right) = -\frac{1}{\sqrt{y}} \frac{\partial u}{\partial \sigma} + 4y \frac{\partial^2 u}{\partial \sigma^2}$$

带入原方程可得

$$4y \frac{\partial^2 u}{\partial \rho^2} - \frac{1}{\sqrt{y}} \frac{\partial u}{\partial \sigma} + 4y \frac{\partial^2 u}{\partial \sigma^2} = 0 \implies \frac{\partial^2 u}{\partial \rho^2} + \frac{\partial^2 u}{\partial \sigma^2} + \frac{1}{3\sigma} \frac{\partial u}{\partial \sigma} = 0. \square$$

4. 见书, 略.

5. 利用分离变量法分离下列方程

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = 0$$

解: 该方程为三元, 则设  $u(r, \theta, z) = v(r, \theta) Z(z)$ , 则有

$$\frac{1}{r} \frac{\partial}{\partial r} \left[ Z(z) r \frac{\partial v}{\partial r} \right] + Z(z) \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2} + v(r, \theta) \frac{d^2 Z(z)}{dz^2} = 0$$

分离可得

$$\frac{\frac{1}{r} \frac{\partial}{\partial r} \left[ r \frac{\partial v}{\partial r} \right] + \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2}}{v(r, \theta)} = -\frac{\frac{d^2 Z(z)}{dz^2}}{Z(z)} = -\lambda$$

即

$$\begin{cases} \frac{1}{r} \frac{\partial}{\partial r} \left[ r \frac{\partial v}{\partial r} \right] + \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2} + \lambda v(r, \theta) = 0 \\ \frac{d^2 Z(z)}{dz^2} - \lambda Z(z) = 0 \end{cases}$$

对于第一个方程，继续利用分离变量法，设  $v(r, \theta) = R(r)\Theta(\theta)$ ，则有

$$\frac{1}{r} \frac{d}{dr} \left[ \Theta(\theta) r \frac{dR(r)}{dr} \right] + R(r) \frac{1}{r^2} \frac{d^2 \Theta(\theta)}{d\theta^2} + \lambda R(r) \Theta(\theta) = 0$$

分离可得

$$\frac{\frac{1}{r} \frac{d}{dr} \left[ r \frac{dR(r)}{dr} \right] + \lambda R(r)}{R(r)} = - \frac{\frac{d^2 \Theta(\theta)}{d\theta^2}}{\Theta(\theta)} = -\mu$$

即

$$\begin{cases} \frac{1}{r} \frac{d}{dr} \left[ r \frac{dR(r)}{dr} \right] + \lambda R(r) + \mu \frac{R(r)}{r^2} = 0 \\ \frac{d^2 \Theta(\theta)}{d\theta^2} - \mu \Theta(\theta) = 0 \end{cases}$$

综上可得

$$\begin{cases} \frac{1}{r} \frac{d}{dr} \left[ r \frac{dR(r)}{dr} \right] + \left( \lambda + \frac{\mu}{r^2} \right) R(r) = 0 \\ \frac{d^2 \Theta(\theta)}{d\theta^2} - \mu \Theta(\theta) = 0 \\ \frac{d^2 Z(z)}{dz^2} - \lambda Z(z) = 0 \end{cases}$$

## 6. Fourier 推导无界弦上的自由振动

$$\frac{\partial^2 u}{\partial t^2} - a^2 \frac{\partial^2 u}{\partial x^2} = 0$$

$$u|_{t=0} = \phi(x), \quad \frac{\partial u}{\partial t} \Big|_{t=0} = \psi(x)$$

解：作 Fourier 变换有

$$U(k, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u(x, t) e^{-ikx} dx, \quad \Phi(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \phi(x) e^{-ikx} dx, \quad \Psi(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x) e^{-ikx} dx$$

且有

$$\mathcal{F} \left[ \frac{\partial^2 u}{\partial x^2} \right] = (ik)^2 \mathcal{F}[u(x, t)] = -k^2 U(k, t)$$

则可得

$$\frac{d^2 U(k, t)}{dt^2} + k^2 a^2 U(k, t) = 0$$

$$U(k, t)|_{t=0} = \Phi(k), \quad \frac{\partial U(k, t)}{\partial t} \Big|_{t=0} = \Psi(k)$$

解得通解为

$$U(k, t) = A \sin k a t + B \cos k a t$$

带入边界条件可得

$$A = \frac{\Psi(k)}{ka} \quad B = \Phi(k)$$

则有

$$U(k, t) = \Phi(k) \cos k a t + \frac{\Psi(k)}{ka} \sin k a t$$

利用 Fourier 反演公式可得

$$u(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} [\Phi(k) \cos k a t + \frac{\Psi(k)}{ka} \sin k a t] e^{ikx} dk$$

对于  $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Phi(k) \cos k a t \cdot e^{ikx} dk$  有

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Phi(k) \cos k a t \cdot e^{ikx} dk &= \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{2} \int_{-\infty}^{\infty} \Phi(k) [e^{ikat} + e^{-ikat}] \cdot e^{ikx} dk \\ &= \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{2} \int_{-\infty}^{\infty} \Phi(k) [e^{ik(x+at)} + e^{ik(x-at)}] dk \\ &= \frac{1}{2} \left\{ \frac{1}{\sqrt{2\pi}} \left[ \int_{-\infty}^{\infty} \Phi(k) e^{ik(x+at)} dk + \int_{-\infty}^{\infty} \Phi(k) e^{ik(x-at)} dk \right] \right\} \\ &= \frac{1}{2} [\phi(x+at) + \phi(x-at)] \end{aligned}$$

对于  $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\Psi(k)}{ka} \sin k a t \cdot e^{ikx} dk$  有

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\Psi(k)}{ka} \sin k a t \cdot e^{ikx} dk &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(k) \left[ \int_0^t \cos k a \tau d\tau \right] \cdot e^{ikx} dk \\ &= \int_0^t \left[ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(k) \cos k a \tau \cdot e^{ikx} dk \right] d\tau \\ &= \frac{1}{2} \int_0^t [\psi(x+at) + \psi(x-at)] d\tau \\ &= \frac{1}{2a} \int_{x-at}^{x+at} \psi(\xi) d\xi \end{aligned}$$

综上可得

$$u(x, t) = \frac{1}{2} [\phi(x+at) + \phi(x-at)] + \frac{1}{2a} \int_{x-at}^{x+at} \psi(\xi) d\xi. \square$$

7. Poisson 第一类边值问题 Green 函数，导出下列定解问题的解

$$\begin{cases} \nabla^2 u(\mathbf{r}) = -f(\mathbf{r}) \\ u(\mathbf{r})|_{\Sigma} = h(\mathbf{r}) \end{cases}$$

解：其对应的 Green 函数为

$$\begin{cases} \nabla^2 G(\mathbf{r}; \mathbf{r}') = -\frac{1}{\alpha} \delta(\mathbf{r} - \mathbf{r}') \\ G(\mathbf{r}; \mathbf{r}')|_{\Sigma} = 0 \end{cases}$$

第一个方程乘  $G(\mathbf{r}; \mathbf{r}')$  减去第二个方程乘  $u(\mathbf{r})$  可得

$$G(\mathbf{r}; \mathbf{r}') \nabla^2 u(\mathbf{r}) - u(\mathbf{r}) \nabla^2 G(\mathbf{r}; \mathbf{r}') = - \left[ G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) - \frac{1}{\alpha} u(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') \right]$$

积分可得

$$\begin{aligned} \iiint_V [G(\mathbf{r}; \mathbf{r}') \nabla^2 u(\mathbf{r}) - u(\mathbf{r}) \nabla^2 G(\mathbf{r}; \mathbf{r}')] dV &= - \iiint_V \left[ G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) - \frac{1}{\alpha} u(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') \right] dV \\ \iint_{\Sigma} [G(\mathbf{r}; \mathbf{r}') \nabla u(\mathbf{r}) - u(\mathbf{r}) \nabla G(\mathbf{r}; \mathbf{r}')] d\Sigma &= \frac{1}{\alpha} u(\mathbf{r}') - \iiint_V G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) dV \end{aligned}$$

则有

$$\begin{aligned} u(\mathbf{r}') &= \alpha \iiint_V G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) dV + \alpha \iint_{\Sigma} [\textcolor{red}{G(\mathbf{r}; \mathbf{r}')}|_{\Sigma} \nabla u(\mathbf{r})|_{\Sigma} - u(\mathbf{r})|_{\Sigma} \nabla G(\mathbf{r}; \mathbf{r}')|_{\Sigma}] d\Sigma \\ &= \alpha \iiint_V G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) dV - \alpha \iint_{\Sigma} \textcolor{red}{u(\mathbf{r})}|_{\Sigma} \nabla \textcolor{blue}{G(\mathbf{r}; \mathbf{r}')}|_{\Sigma} d\Sigma \\ &= \alpha \iiint_V G(\mathbf{r}; \mathbf{r}') f(\mathbf{r}) dV - \alpha \iint_{\Sigma} \textcolor{red}{h(\mathbf{r})} \frac{\partial G(\mathbf{r}; \mathbf{r}')}{\partial \mathbf{n}} d\Sigma \end{aligned}$$

本资源免费共享 收集网站 nuaa.store

将  $\mathbf{r}$  和  $\mathbf{r}'$  对调一下可得

$$u(\mathbf{r}) = \alpha \iiint_V G(\mathbf{r}'; \mathbf{r}) f(\mathbf{r}') dV' - \alpha \iint_{\Sigma} h(\mathbf{r}') \frac{\partial G(\mathbf{r}'; \mathbf{r})}{\partial \mathbf{n}'} d\Sigma'. \square$$

## 第二部分 2023 年部分作业题解析

## 1. (P170) 第十章 T2.(1)

解：直接写出通解表达式为

$$g(t; \tau) = \begin{cases} A(\tau) \sinh kt + B(\tau) \cosh kt, & t < \tau \\ C(\tau) \sinh kt + D(\tau) \cosh kt, & t > \tau \end{cases}$$

带入边界条件可得

$$A(\tau) = 0$$

$$B(\tau) = 0$$

$$C(\tau) \sinh k\tau + D(\tau) \cosh k\tau = 0$$

$$C(\tau) \cosh k\tau + D(\tau) \sinh k\tau = \frac{1}{k}$$

利用 Cramer 法则解之可得

$$C(\tau) = \frac{\begin{vmatrix} 0 & \cosh k\tau \\ \frac{1}{k} & \sinh k\tau \end{vmatrix}}{\begin{vmatrix} \sinh k\tau & \cosh k\tau \\ \cosh k\tau & \sinh k\tau \end{vmatrix}} = \frac{1}{k} \cosh k\tau \quad D(\tau) = \frac{\begin{vmatrix} \sinh k\tau & 0 \\ \cosh k\tau & \frac{1}{k} \end{vmatrix}}{\begin{vmatrix} \sinh k\tau & \cosh k\tau \\ \cosh k\tau & \sinh k\tau \end{vmatrix}} = -\frac{1}{k} \sinh k\tau$$

则有

$$g(t; \tau) = \frac{1}{k} \sinh k(t - \tau), \quad t > \tau$$

综上可得

$$g(t; \tau) = \frac{1}{k} \sinh k(t - \tau) \eta(t - \tau), \quad t, \tau > 0$$

## 2. (P171) 第十章 T3

解：其对应的格林函数为

$$\frac{d^2 g(t; \tau)}{d\tau^2} + k^2 g(t; \tau) = \delta(t - \tau), \quad t, \tau > 0, \quad k > 0$$

$$g(t; \tau)|_{\tau > t} = 0, \quad \left. \frac{dg(t; \tau)}{dt} \right|_{\tau > t} = 0$$

原条件变为

$$\frac{d^2 y(\tau)}{d\tau^2} + k^2 y(\tau) = f(\tau), \quad \tau > 0, \quad k > 0$$

$$y(0) = A, \quad \left. \frac{dy(\tau)}{d\tau} \right|_{\tau=0} = B$$

做差可得

$$g(t; \tau) \frac{d^2 y(\tau)}{d\tau^2} - y(\tau) \frac{d^2 g(t; \tau)}{d\tau^2} = g(t; \tau) f(\tau) - y(\tau) \delta(t - \tau)$$

积分可得

$$\int_0^\infty \left[ g(t; \tau) \frac{d^2 y(\tau)}{d\tau^2} - y(\tau) \frac{d^2 g(t; \tau)}{d\tau^2} \right] d\tau = \int_0^\infty g(t; \tau) f(\tau) d\tau - \int_0^\infty y(\tau) \delta(t - \tau) d\tau$$

$$\left. \left[ g(t; \tau) \frac{dy(\tau)}{d\tau} - y(\tau) \frac{dg(t; \tau)}{d\tau} \right] \right|_{\tau=0} = \int_0^t g(t; \tau) f(\tau) d\tau - y(t)$$

其中蕴含了  $g(t; \tau)|_{\tau>t} = 0$ ,  $\left. \frac{dg(t; \tau)}{dt} \right|_{\tau>t} = 0$ , 解之可得

$$y(t) = \int_0^t g(t; \tau) f(\tau) d\tau + \left. \left[ g(t; \tau) \frac{dy(\tau)}{d\tau} - y(\tau) \frac{dg(t; \tau)}{d\tau} \right] \right|_{\tau=0}$$

$$= \int_0^t g(t; \tau) f(\tau) d\tau + \left. \left[ Bg(t; \tau) - A \frac{dg(t; \tau)}{d\tau} \right] \right|_{\tau=0}$$

因为 Green 函数的通解为  $g(t; \tau) = \frac{1}{k} \sin k(t - \tau)$ , 带入可得

$$y(t) = A \cos kt + \frac{B}{k} \sin kt + \frac{1}{k} \int_0^t \sin k(t - \tau) f(\tau) d\tau. \square$$

### 3. (P171)第十章 T4.(1)

解: 步骤同上

因为 Green 函数的通解为  $g(t; \tau) = \frac{1}{k} \sinh k(t - \tau)$ , 带入可得

$$y(t) = A \cosh kt + \frac{B}{k} \sinh kt + \frac{1}{k} \int_0^t \sinh k(t - \tau) f(\tau) d\tau. \square$$

注: 在这两个例子中, 若  $y(0) = 0$ ,  $\left. \frac{dy(\tau)}{d\tau} \right|_{\tau=0} = 0$ , 则分别有

$$(1) y(t) = \frac{1}{k} \int_0^t \sinh k(t - \tau) f(\tau) d\tau$$

$$(2) y(t) = \frac{1}{k} \int_0^t \sinh k(t - \tau) f(\tau) d\tau$$

也就是说

$$\frac{d^2 y(t)}{dt^2} + k^2 y(t) = f(t), t > 0, k > 0$$

$$y(0) = 0, \left. \frac{dy(t)}{dt} \right|_{t=0} = 0$$

解为

$$y(t) = \frac{1}{k} \int_0^t \sinh k(t - \tau) f(\tau) d\tau. \square$$

$$\frac{d^2 y(t)}{dt^2} - k^2 y(t) = f(t), t > 0, k > 0$$

$$y(0) = 0, \left. \frac{dy(t)}{dt} \right|_{t=0} = 0$$

解为

$$y(t) = \frac{1}{k} \int_0^t \sinh k(t-\tau) f(\tau) d\tau. \square$$

## 4. (P171)第十章 T5.(1)

解：直接写出通解为

$$g(x; \xi) = \begin{cases} A(\xi) \sinh kx + B(\xi) \cosh kx, & x < \xi \\ C(\xi) \sinh kx + D(\xi) \cosh kx, & x > \xi \end{cases}$$

带入边界条件可得

$$B(\xi) = 0 \quad C(\xi) \cosh k\xi + D(\xi) \sinh k\xi - A(\xi) \cosh k\xi = \frac{1}{k}$$

$$C(\xi) \sinh k + D(\xi) \cosh k = 0 \quad A(\xi) \sinh k\xi = C(\xi) \sinh k\xi + D(\xi) \cosh k\xi$$

利用 Cramer 法则解之可得

$$\begin{aligned} A(\xi) &= -\frac{1}{k} \frac{\sinh k(1-\xi)}{\sinh k} & B(\xi) &= 0 \\ C(\xi) &= \frac{1}{k} \frac{\cosh k \sinh k \xi}{\sinh k} & D(\xi) &= -\frac{1}{k} \sinh k \xi \end{aligned}$$

则有

$$g(x; \xi) = \begin{cases} -\frac{1}{k} \frac{\sinh k(1-\xi)}{\sinh k} \cdot \sinh kx, & x < \xi \\ \frac{1}{k} \frac{\sinh k \xi \sinh k(1-x)}{\sinh k}, & x > \xi \end{cases}$$

合并为

$$g(x; \xi) = -\frac{1}{k} \frac{\sinh k(1-\xi)}{\sinh k} \cdot \sinh kx - \frac{1}{k} \left[ \frac{\sinh k \xi \sinh k(1-x)}{\sinh k} - \frac{\sinh k(1-\xi)}{\sinh k} \cdot \sinh kx \right] \eta(x-\xi)$$

最终化简可得

$$g(x; \xi) = -\frac{1}{k} \frac{\sinh k(1-\xi)}{\sinh k} \cdot \sinh kx + \frac{1}{k} \sinh k(x-\xi) \eta(x-\xi). \square$$

注：部分化简过程如下

$$\begin{aligned} & \frac{\sinh k \xi \sinh k(1-x)}{\sinh k} - \frac{\sinh k(1-\xi)}{\sinh k} \cdot \sinh kx = \frac{\sinh k \xi \sinh k(1-x) - \sinh kx \sinh k(1-\xi)}{\sinh k} \\ &= \frac{\sinh k \cosh kx \sinh k \xi - \cosh k \sinh kx \sinh k \xi - \sinh k \sinh kx \cosh k \xi + \cosh k \sinh kx \sinh k \xi}{\sinh k} \\ &= -\sinh k(x-\xi) \end{aligned}$$

## 5. (P225)第十三章 T5

解：利用分离变量法可得

$$\begin{cases} X''(x) - \lambda X(x) = 0 \\ Y''(y) + \lambda Y(y) = 0 \end{cases}$$

则有

$$Y(y) = A \sin \sqrt{\lambda} y + B \cos \sqrt{\lambda} y, \quad Y'(y) = \sqrt{\lambda} (A \cos \sqrt{\lambda} y - B \sin \sqrt{\lambda} y)$$

带入边界条件可得

$$A=0 \quad B=1 \quad \lambda_n = \left(\frac{n\pi}{b}\right)^2$$

可得

$$Y_n(y) = \cos \frac{n\pi}{b} y$$

组装可得

$$u(x,y) = Cx + D + \sum_{n=1}^{\infty} \left( C_n \sinh \frac{n\pi}{b} x + D_n \cosh \frac{n\pi}{b} x \right) \cos \frac{n\pi}{b} y$$

带入边界条件可得

$$\begin{aligned} u(0,y) &= D + \sum_{n=1}^{\infty} D_n \cos \frac{n\pi}{b} y = u_0 \\ u(a,y) &= Ca + D + \sum_{n=1}^{\infty} \left( C_n \sinh \frac{n\pi a}{b} + D_n \cosh \frac{n\pi a}{b} \right) \cos \frac{n\pi}{b} y = u_0 \left[ 3\left(\frac{y}{b}\right)^2 - 2\left(\frac{y}{b}\right)^3 \right] \end{aligned}$$

则有

$$D = u_0 \quad D_n = 0$$

$$C_n = - \frac{48u_0}{(2n+1)^4 \pi^4 \sinh \frac{(2n+1)\pi a}{b}}$$

再由  $u(a,b) = u_0$  可得

$$C = - \frac{u_0}{2a}$$

综上可得

$$u(x,y) = u_0 \left( 1 - \frac{x}{2a} \right) - \frac{48u_0}{\pi^4} \sum_{n=0}^{\infty} \frac{\sinh \frac{(2n+1)\pi}{b} x}{(2n+1)^4 \sinh \frac{(2n+1)\pi a}{b}} \cos \frac{(2n+1)\pi}{b} y. \square$$

注：部分计算过程如下

$$D_n = \frac{2}{b} \int_0^b (u_0 - D) \cos \frac{n\pi}{b} y dy = \frac{2(u_0 - D)}{n\pi} \int_0^{n\pi} \cos u du = 0$$

$$\begin{aligned}
C_n &= \frac{2}{b \sinh \frac{n\pi a}{b}} \int_0^b \left\{ u_0 \left[ 3 \left( \frac{y}{b} \right)^2 - 2 \left( \frac{y}{b} \right)^3 \right] - u_0 - Ca \right\} \cos \frac{n\pi}{b} y \, dy \\
&= \frac{2}{b \sinh \frac{n\pi a}{b}} \left[ 3u_0 \int_0^b \left( \frac{y}{b} \right)^2 \cos \frac{n\pi}{b} y \, dy - 2u_0 \int_0^b \left( \frac{y}{b} \right)^3 \cos \frac{n\pi}{b} y \, dy - (u_0 + Ca) \int_0^b \cos \frac{n\pi}{b} y \, dy \right] \\
&= \frac{2}{b \sinh \frac{n\pi a}{b}} \left[ 3u_0 b \int_0^1 y^2 \cos n\pi y \, dy - 2u_0 b \int_0^1 y^3 \cos n\pi y \, dy - (u_0 + Ca) b \int_0^1 \cos n\pi y \, dy \right] \\
&= \frac{2}{b \sinh \frac{n\pi a}{b}} \left[ 3u_0 b \cdot \frac{2n\pi \cos n\pi}{n^3 \pi^3} - 2u_0 b \cdot \frac{3n^2 \pi^2 \cos n\pi + 6(1 - \cos n\pi)}{n^4 \pi^4} \right] \\
&= -\frac{2u_0}{\sinh \frac{n\pi a}{b}} \frac{12(1 - \cos n\pi)}{n^4 \pi^4} = -\frac{24u_0(1 - \cos n\pi)}{n^4 \pi^4 \sinh \frac{n\pi a}{b}} = -\frac{48u_0}{(2n+1)^4 \pi^4 \sinh \frac{(2n+1)\pi a}{b}} \\
&\quad \sum_{n=0}^{\infty} \frac{1}{(2n+1)^4} = \frac{\pi^4}{96}
\end{aligned}$$

6. (P226)第十三章 T7.(1)

解：设  $u(x, y) = v(x) + w(x, y)$ ，则有

$$\begin{aligned}
\nabla^2 v(x) &= -2 \\
v|_{x=0,a} &= 0
\end{aligned}$$

解之可得

$$v(x) = x(a-x)$$

而又有

$$\begin{aligned}
\nabla^2 w(x, y) &= 0 \\
w|_{x=0,a} &= -v|_{x=0,a} = 0, \quad w|_{y=\pm\frac{b}{2}} = -v|_{y=\pm\frac{b}{2}} = -x(a-x)
\end{aligned}$$

带入一般解有

$$w(x, y) = \sum_{n=1}^{\infty} \left( C_n \sinh \frac{n\pi}{a} y + D_n \cosh \frac{n\pi}{a} y \right) \sin \frac{n\pi}{a} x$$

带入边界条件可得

$$\begin{aligned}
C_n &= 0 \\
D_n &= -\frac{8a^2}{(2n+1)^3 \pi^3 \cosh \frac{2n+1}{2a} \pi b}
\end{aligned}$$

则有

$$w(x, y) = -\frac{8a^2}{\pi^3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^3} \frac{\cosh \frac{2n+1}{a} \pi y}{\cosh \frac{2n+1}{2a} \pi b} \sin \frac{2n+1}{a} \pi x$$

综上可得

$$u(x,y) = x(a-x) - \frac{8a^2}{\pi^3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^3} \frac{\cosh \frac{2n+1}{a}\pi y}{\cosh \frac{2n+1}{2a}\pi b} \sin \frac{2n+1}{a}\pi x. \square$$

注：部分计算过程如下

$$w(x,y) = w(x, -y) \implies C_n = 0$$

$$\begin{aligned} D_n &= -\frac{2}{a \cosh \frac{n}{2a}\pi b} \int_0^a x(a-x) \sin \frac{n}{a}\pi x dx = -\frac{2}{a \cdot \cosh \frac{n}{2a}\pi b} \cdot \frac{2a^3(\cos n\pi - 1)}{n^3\pi^3} \\ &= -\frac{2}{a \cdot \cosh \frac{2n+1}{2a}\pi b} \cdot \frac{4a^3}{(2n+1)^3\pi^3} = -\frac{8a^2}{(2n+1)^3\pi^3 \cosh \frac{2n+1}{2a}\pi b} \end{aligned}$$

7. (P226)第十三章 T7.(2)

解：步骤同上，解得

$$v(x,y) = \frac{1}{12} xy(a^3 - x^3)$$

且有

$$\begin{aligned} \nabla^2 w(x,y) &= 0 \\ w|_{x=0,a} &= -v|_{x=0,a} = 0, \quad w|_{y=\pm\frac{b}{2}} = -v|_{y=\pm\frac{b}{2}} = \pm \frac{b}{24} x(a^3 - x^3) \end{aligned}$$

解之可得 资源免费共享 收集网站 nuaa.store

$$w(x,y) = \frac{a^4 b}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} \frac{\sinh \frac{n\pi}{a} y}{\sinh \frac{n\pi}{2a} b} \sin \frac{n\pi}{a} x + \frac{4a^4 b}{\pi^5} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^5} \frac{\sinh \frac{2n+1}{a}\pi y}{\sinh \frac{2n+1}{2a}\pi b} \sin \frac{2n+1}{a}\pi x$$

综上可得

$$\begin{aligned} u(x,y) &= \frac{1}{12} xy(a^3 - x^3) + \frac{a^4 b}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} \frac{\sinh \frac{n\pi}{a} y}{\sinh \frac{n\pi}{2a} b} \sin \frac{n\pi}{a} x \\ &\quad + \frac{4a^4 b}{\pi^5} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^5} \frac{\sinh \frac{2n+1}{a}\pi y}{\sinh \frac{2n+1}{2a}\pi b} \sin \frac{2n+1}{a}\pi x. \square \end{aligned}$$

注：部分计算过程如下

$$w(x,y) = -w(x, -y) \implies D_n = 0$$

$$\begin{aligned}
C_n &= \frac{2}{a \sinh \frac{n}{2a} \pi b} \int_0^a \frac{b}{24} x(a^3 - x^3) \sin \frac{n}{a} \pi x \, dx = \frac{a^2 b}{12} \int_0^a x \sin \frac{n}{a} \pi x \, dx - \frac{b}{12a} \int_0^a x^4 \sin \frac{n}{a} \pi x \, dx \\
&= \frac{a^2 b}{12 \sinh \frac{n}{2a} \pi b} \cdot \frac{a^2 \cos n \pi}{n \pi} - \frac{2}{a \cdot \sinh \frac{n}{2a} \pi b} \cdot \frac{b}{24} \cdot \frac{a^5 (n^4 \pi^4 \cos n \pi - 12n^2 \pi^2 \cos n \pi + 24 \cos n \pi - 24)}{n^5 \pi^5} \\
&= \frac{a^4 b (-1)^n}{12 \pi n \cdot \sinh \frac{n}{2a} \pi b} - \frac{b}{12a \cdot \sinh \frac{n}{2a} \pi b} \cdot \left[ \frac{a^5 (n^4 \pi^4 - 12n^2 \pi^2)}{n^5 \pi^5} \cos n \pi + \frac{24a^5 (\cos n \pi - 1)}{n^5 \pi^5} \right] \\
&= \frac{a^4 b (-1)^n}{12 \pi n \cdot \sinh \frac{n}{2a} \pi b} - \frac{b}{12 \sinh \frac{n}{2a} \pi b} \cdot \frac{a^4 (n^4 \pi^4 - 12n^2 \pi^2)}{n^5 \pi^5} \cdot (-1)^n + \frac{b}{\sinh \frac{n}{2a} \pi b} \cdot \frac{2a^4 (\cos n \pi - 1)}{n^5 \pi^5} \\
&= - \left[ \frac{a^4 b}{12 \pi} \frac{1}{n \sinh \frac{n}{2a} \pi b} + \frac{a^4 b}{12 \pi} \frac{1}{n \cdot \sinh \frac{n}{2a} \pi b} - \frac{a^4 b}{\pi^3} \frac{1}{n^3 \cdot \sinh \frac{n}{2a} \pi b} \right] (-1)^n + \frac{4a^4 b}{\pi^5} \cdot \frac{1}{(2n+1)^5 \sinh \frac{2n+1}{2a} \pi b} \\
&= \frac{a^4 b}{\pi^3} \frac{(-1)^n}{n^3 \cdot \sinh \frac{n}{2a} \pi b} + \frac{4a^4 b}{\pi^5} \cdot \frac{1}{(2n+1)^5 \sinh \frac{2n+1}{2a} \pi b}
\end{aligned}$$

## 8. (P341)第十八章 T5

解：作 Fourier 变换可得

$$\begin{aligned}
U(k, t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u(x, t) e^{-ikx} \, dx, \quad \Phi(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \phi(x) e^{-ikx} \, dx \\
\Psi(k) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x) e^{-ikx} \, dx, \quad F(k, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x, t) e^{-ikx} \, dx
\end{aligned}$$

方程变形为

$$\begin{aligned}
\frac{d^2 U(k, t)}{dt^2} + k^2 a^2 U(k, t) &= F(k, t) \\
U(k, t)|_{t=0} &= \Phi(k), \quad \left. \frac{\partial U(k, t)}{\partial t} \right|_{t=0} = \Psi(k)
\end{aligned}$$

设  $U(k, t) = v(k, t) + w(k, t)$ , 则有

$$\begin{aligned}
\frac{d^2 v(k, t)}{dt^2} + k^2 a^2 v(k, t) &= F(k, t) \\
v(k, t)|_{t=0} &= 0, \quad \left. \frac{\partial v(k, t)}{\partial t} \right|_{t=0} = 0
\end{aligned}$$

解之可得(见第二部分 3)

$$v(k, t) = \frac{1}{ka} \int_0^t \sin ka(t - \tau) f(k, \tau) \, d\tau$$

且有

$$\frac{d^2 w(k,t)}{dt^2} + k^2 a^2 w(k,t) = 0$$

$$w(k,t)|_{t=0} = \Phi(k), \quad \left. \frac{\partial w(k,t)}{\partial t} \right|_{t=0} = \Psi(k)$$

解之可得(见第一部分 6)

$$w(k,t) = \frac{\Psi(k)}{ka} \sin kat + \Phi(k) \cos kat$$

可得

$$U(k,t) = \frac{\Psi(k)}{ka} \sin kat + \Phi(k) \cos kat + \frac{1}{ka} \int_0^t \sin ka(t-\tau) f(k,\tau) d\tau$$

利用 Fourier 反演公式可得

$$u(x,t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \left[ \frac{\Psi(k)}{ka} \sin kat + \Phi(k) \cos kat + \frac{1}{ka} \int_0^t \sin ka(t-\tau) f(k,\tau) d\tau \right] e^{ikx} dk$$

前两项解之可得

$$\frac{1}{2} [\phi(x+at) + \phi(x-at)] + \frac{1}{2a} \int_{x-at}^{x+at} \psi(\xi) d\xi$$

对于  $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{ka} \int_0^t \sin ka(t-\tau) f(k,\tau) d\tau e^{ikx} dk$  有

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{ka} \int_0^t \sin ka(t-\tau) f(k,\tau) d\tau e^{ikx} dk &= \int_0^t \left[ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(k,\tau) \frac{\sin ka(t-\tau)}{ka} e^{ikx} dk \right] d\tau \\ &= \int_0^t \left[ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(k,\tau) \left[ \int_0^\tau \cos ka(t-s) ds \right] \cdot e^{ikx} dk \right] d\tau \\ &= \int_0^t \left\{ \int_0^\tau \left[ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(k,\tau) \cos ka(t-s) e^{ikx} dk \right] ds \right\} d\tau \\ &= \int_0^t \left\{ \frac{1}{2} \int_0^\tau [f(x+a(t-s),\tau) + f(x-a(t-s),\tau)] ds \right\} d\tau \\ &= \frac{1}{2a} \int_0^t \left[ \int_{x-a(t-\tau)}^{x+a(t-\tau)} f(\xi,\tau) d\xi \right] d\tau \end{aligned}$$

综上可得

$$u(x,t) = \frac{1}{2} [\phi(x+at) + \phi(x-at)] + \frac{1}{2a} \int_{x-at}^{x+at} \psi(\xi) d\xi + \frac{1}{2a} \int_0^t \left[ \int_{x-a(t-\tau)}^{x+a(t-\tau)} f(\xi,\tau) d\xi \right] d\tau. \square$$

8. 求解下列非齐次方程的定解问题

$$\nabla^2 u = -2Ax$$

$$u(0,y) = 0, \quad u(a,y) = 0, \quad u(x,0) = 0, \quad u(x,b) = 0$$

解：设  $u(x,y) = v(x) + w(x,y)$ ，则有

$$\begin{aligned}\nabla^2 v(x) &= -2Ax \\ v(0,y) &= 0, \quad v(a,y) = 0\end{aligned}$$

解之可得

$$v(x) = -\frac{1}{3}Ax(x^2 - a^2)$$

且有

$$\begin{aligned}\nabla^2 w(x,y) &= 0 \\ w(0,y) &= 0, \quad w(a,y) = 0, \quad w(x,0) = -v(x,0), \quad w(x,b) = -v(x,b)\end{aligned}$$

带入一般解可得

$$w(x,y) = \sum_{n=1}^{\infty} \left( C_n \sinh \frac{n\pi}{a} y + D_n \cosh \frac{n\pi}{a} y \right) \sin \frac{n\pi}{a} x$$

带入边界条件有

$$\begin{aligned}w(x,0) &= \sum_{n=1}^{\infty} D_n \sin \frac{n\pi}{a} x = \frac{1}{3}Ax(x^2 - a^2) \\ w(x,b) &= \sum_{n=1}^{\infty} \left( C_n \sinh \frac{n\pi b}{a} + D_n \cosh \frac{n\pi b}{a} \right) \sin \frac{n\pi}{a} x = \frac{1}{3}Ax(x^2 - a^2)\end{aligned}$$

即

$$\begin{aligned}D_n &= \frac{2A}{3a} \int_0^a x(x^2 - a^2) \sin \frac{n\pi}{a} x \, dx \\ C_n \sinh \frac{n\pi b}{a} + D_n \cosh \frac{n\pi b}{a} &= \frac{2A}{3a} \int_0^a x(x^2 - a^2) \sin \frac{n\pi}{a} x \, dx = D_n\end{aligned}$$

解得

$$D_n = -\frac{4Aa^3(-1)^n}{n^3\pi^3} \quad C_n = \frac{1 - \cosh \frac{n\pi b}{a}}{\sinh \frac{n\pi b}{a}} D_n = -\frac{4Aa^3(-1)^n}{n^3\pi^3} \cdot \frac{1 - \cosh \frac{n\pi b}{a}}{\sinh \frac{n\pi b}{a}}$$

则有

$$\begin{aligned}w(x,y) &= -\frac{4Aa^3}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} \left[ \frac{1 - \cosh \frac{n\pi b}{a}}{\sinh \frac{n\pi b}{a}} \sinh \frac{n\pi}{a} y + \cosh \frac{n\pi}{a} y \right] \sin \frac{n\pi}{a} x \\ &= -\frac{8Aa^3}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 \sinh \frac{n\pi b}{a}} \sinh \frac{n\pi b}{2a} \cosh \frac{n\pi b}{2a} \sin \frac{n\pi}{a} x\end{aligned}$$

综上可得

$$u(x,y) = -\frac{1}{3}Ax(x^2 - a^2) - \frac{8Aa^3}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 \sinh \frac{n\pi b}{a}} \sinh \frac{n\pi b}{2a} \cosh \frac{n\pi b}{2a} \sin \frac{n\pi}{a} x. \square$$

本资源免费共享 收集网站 nuaa.store

## 第三部分 部分必备结论

$$1. \int_{-\infty}^{\infty} e^{-t^2} \cos 2zt dt = \sqrt{\pi} e^{-z^2}.$$

$$2. \int_0^{n\pi} x \sin x dx = -n\pi \cos n\pi, \quad \int_0^{n\pi} x \cos x dx = \cos n\pi - 1.$$

$$3. \int_0^{n\pi} x^2 \sin x dx = 2(\cos n\pi - 1) - n^2 \pi^2 \cos n\pi, \quad \int_0^{n\pi} x^2 \cos x dx = 2n\pi \cos n\pi.$$

$$4. u(x,t)|_{x=0} = u_0 \xrightarrow{\text{Laplace}} U(x,p)|_{x=0} = \frac{u_0}{p}.$$

$$5. (1) b^2 - ac < 0, \rho = \xi + \eta, \sigma = i(\xi - \eta); (2) b^2 - ac > 0, \rho = \xi + \eta, \sigma = \xi - \eta.$$

$$6. f^{(n)}(t) \doteq p^n F(p) - \sum_{k=0}^{n-1} p^k f^{(n-1-k)}(0).$$

$$7. \mathcal{F}\{f^{(n)}(t)\} = (ik)^n \mathcal{F}\{f(t)\}.$$

$$8. \mathcal{F}[f_1(t)] \mathcal{F}[f_2(t)] = \mathcal{F}\left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_1(\xi) f_2(t-\xi) d\xi\right].$$

$$9. \delta(at - |x - x'|) = \begin{cases} 0, & |x - x'| \neq at \\ \infty, & |x - x'| = at \end{cases}.$$

$$10. \eta(at - |x - x'|) = \begin{cases} 0, & |x - x'| > at \\ 1, & |x - x'| < at \end{cases}.$$

$$11. \mathcal{L}(1) = \frac{1}{p}.$$

$$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)], \cos \alpha \sin \beta = \frac{1}{2} [\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

$$12. \cos \alpha \cos \beta = \frac{1}{2} [\cos(\alpha + \beta) + \cos(\alpha - \beta)], \sin \alpha \sin \beta = -\frac{1}{2} [\cos(\alpha + \beta) - \cos(\alpha - \beta)].$$

$$13. \frac{dy}{dx} + p(x)y = f(x) \implies u = \int p(x) dx, y = e^{-u} \left[ C + \int f(x) e^u dx \right].$$

$$14. \text{Green 函数在 } x = \xi \text{ 处连续, 其对时间的导数在 } x = \xi \text{ 处有一个跃度 } \frac{1}{p(t)}.$$

$$15. \frac{d^2 y(t)}{dt^2} - k^2 y(t) = f(t), t > 0, k > 0, y(x)|_{x \rightarrow \pm\infty} \text{ 有界, 则有}$$

$$y(x) = -\frac{1}{2k} \int_{-\infty}^{\infty} e^{-k|x-\xi|} f(\xi) d\xi.$$

16.  $\sinh(x+y) = \sinh x \cosh y + \cosh x \sinh y, \sinh(x-y) = \sinh x \cosh y - \cosh x \sinh y$   
 $\cosh(x+y) = \cosh x \cosh y + \sinh x \sinh y, \cosh(x-y) = \cosh x \cosh y - \sinh x \sinh y$ .
17.  $\cosh^2 x = 1 + \sinh^2 x.$
18.  $e^{-\alpha p} \doteq \delta(t-\alpha).$
19.  $\frac{1}{p} e^{-\alpha p} \doteq \eta(t-\alpha).$
20.  $\frac{1}{p} e^{-\alpha\sqrt{p}} \doteq \operatorname{erfc}\left(\frac{\alpha}{2\sqrt{t}}\right).$
21.  $\frac{1}{\sqrt{p}} e^{-\alpha\sqrt{p}} \doteq \frac{1}{\sqrt{\pi t}} e^{-\frac{\alpha^2}{4t}}.$

本资源免费共享 收集网站 nuaa.store